

On the Consistency of the Ladder Approximation and the Rainbow Approximation of Dyson-Schwinger Equation of QCD

Lei Chang¹, and Yu-xin Liu^{1,2,3,*}

¹ Department of Physics and the MOEC Key Laboratory of Heavy Ion Physics,
Peking University, Beijing 100871, China

² CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China

³ Center of Theoretical Nuclear Physics, National Laboratory of
Heavy Ion Accelerator, Lanzhou 730000, China

February 2, 2008

Abstract

We study the consistency of the ladder approximation and the rainbow approximation of the Dyson-Schwinger equation of QCD. By considering the non-Abelian property of QCD, we show that the QED-type Ward-Takahashi identity is not required for the rainbow-ladder approximation of QCD. It indicates that there does not exist any internal inconsistency in the usual rainbow-ladder approximation of QCD. In addition, we propose a modified ladder approximation which guarantees the Slavnov-Taylor identity for the quark-gluon vertex omitting the ghost effect in the approximation.

PACS numbers: 11.15.Tk, 12.38.Lg, 12.38.Aw

*Corresponding Author

Studies of the truncated sets of Dyson-Schwinger equations have long been pursued as a possible basis for studying nonperturbative aspects of quantum field theories[1]. The simplest truncation scheme is the so-called rainbow-ladder approximation, which has been used extensively and successfully to study various nonperturbative aspects of strong interaction physics (see for example Refs. [2, 4, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15]). Recently, the author of Ref. [16] claims that the ladder approximation to QCD is internally inconsistent with a QED-type Ward-Takahashi relation, then all the results based on the nontrivial solutions of the quark Dyson-Schwinger equation in the ladder approximation should be reconsidered, and its use in the whole energy/momentum range should be abandoned. Since the rainbow-ladder approximation is such a widely used approximation, its internal consistency is then an imperative issue and deserves careful investigation. Then this claim is argued by the authors of Ref. [17], where they pointed out that the Ward-Takahashi relation could not be required in the ladder approximation to QCD and the claim in Ref. [16] is not correct at least in the case of making use of the weak coupling expansion. It is clear that there exists a contradiction between the Ward-Takahashi identity omitting the ghost effect and the ladder approximation in view of the arguments in Ref. [17]. However, an explicit and general verification for the contradiction to be allowed has not yet been given. There leaves then still a room to prove and generalize the reason mentioned in Ref. [17]. One of the aims of this letter is just to give an exact expression for the contradiction and to show that the contradiction is allowed physically to the usual ladder approximation. Then we would propose a modified ladder approximation to conciliate the contradiction. In the following we shall first briefly recall the arguments in Refs. [16] and [17], and then give our statements for the problem.

Let us begin with the rainbow approximation of the quark equation and the ladder truncation of the quark-gluon vertex equation. For a given quark flavor, the quark Dyson-Schwinger equation (DSE) under the rainbow approximation in QCD is expressed [18]

$$S(p)^{-1} = i\gamma \cdot p + m_0 + g^2 C_F \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) \gamma_\mu S(q) \gamma_\nu, \quad (1)$$

where m_0 is the current quark mass, C_F is the eigenvalue of the quadratic Casimir operator in the fundamental representation and $D_{\mu\nu}(k)$ is the dressed-gluon propagator. The DSE of the quark-gluon vertex under the ladder approximation at zero momentum transfer reads

$$\Gamma_\sigma^a(p) = i\gamma_\sigma T^a - g^2 \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) T^b \gamma_\mu S(q) \Gamma_\sigma^a(q) S(q) T^b \gamma_\nu, \quad (2)$$

where a, b are color indices with T^a the standard Gell-Mann $SU(3)$ representation. It is noted that only the one gluon exchange between a quark and an antiquark is allowed in such a ladder approximation, and this is usually referred to as Abelian approximation. It has been shown that, for color singlet vertex (such as vector, axial-vector and pseudoscalar), the

Abelian ladder approximation is valid because of some Ward-Takahashi identities. However, in the case of quark-gluon vertex, the non-Abelian effect would be included in the expression of its equations. There would then exist a contradiction between the Abelian ladder approximation and the Ward-Takahashi identity if one omit the non-Abelian effect. Another fundamental expression is the differential form of Eq.(1), i.e.

$$\frac{\partial S(p)^{-1}}{\partial p_\sigma} = i\gamma_\sigma + g^2 C_F \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) \gamma_\mu \frac{\partial S(q)}{\partial q_\sigma} \gamma_\nu. \quad (3)$$

We also note that this differential form of quark equation is just the special one in the above mentioned ladder approximation. With Eqs.(1), (2) and (3), we can recall the arguments in Refs. [16] and [17].

The author of Ref. [16] argues that, in the ladder approximation, one should omit the ghost-quark scattering kernel contained in the Slavnov-Taylor identity of QCD and therefore the Slavnov-Taylor identity is reduced to the Ward-Takahashi relation

$$\Gamma_\sigma^a(p) = T^a \frac{\partial S^{-1}(p)}{\partial p_\sigma}, \quad (4)$$

which should be valid in the ladder approximation to QCD. Using the above Ward-Takahashi identity and comparing Eqs.(2) and (3), one can obtain a relation

$$-\frac{1}{2}g^2 C_A \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) \gamma_\mu \frac{\partial S(q)}{\partial q_\sigma} \gamma_\nu = 0, \quad (5)$$

where C_A is the eigenvalue of the quadratic Casimir operator in the adjoint representation. From the above relation, the author obtains,

$$\partial_\sigma \Sigma(p) = 0, \quad (6)$$

for the quark self-energy

$$\Sigma(p) = \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) \gamma_\mu S(q) \gamma_\nu, \quad (7)$$

and concludes that, in the ladder approximation of QCD, the quark propagator is only the free one, apart from a redefinition of quark mass, i.e. there is no running/dressed quark mass and in turn the ladder approximation is internally inconsistent.

The authors of Ref. [17] note that Eq. (5) may not be true in the ladder approximation. Based on the weak coupling expansion in Feynman gauge, they obtain mathematically an analytic result, which reads

$$-\frac{1}{2}g^2 C_A \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) \gamma_\mu \frac{\partial S(q)}{\partial q_\sigma} \gamma_\nu \neq 0. \quad (8)$$

Furthermore they argue that Eq. (8) is true, because a lot of numerical calculations with suitable model gluon propagator in specific gauge have given nonzero results. Since the relation in Eq. (5) is directly the result of Eq. (4), the Ward-Takahashi identity should then not be required because the relation in Eq. (5) is not true in weak coupling expansion at least. It is evident that such an argument made in Ref. ([17]) is not solid enough due to a lack of general verification in view of strong interaction physics (QCD). We then try to prove the validity of Eq. (8) in a general point of view.

Due to the property of non-Abelian gauge theory, the gauge fields of QCD are the self-interacting ones. The derivative of the gluon propagator with respect to momentum is related to the three-gluon vertex based on the Slavnov-Taylor identity omitting the ghost effects and can be written as [19]

$$\Gamma_{\mu\nu\sigma}^{3g}(k, k) = -\frac{\partial}{\partial k_\sigma} D_{\mu\nu}^{-1}(k). \quad (9)$$

With such a relation, one has

$$\begin{aligned} & -\frac{1}{2}g^2 C_A \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(p-q) \gamma_\mu \frac{\partial S(q)}{\partial q_\sigma} \gamma_\nu \\ = & \frac{1}{2}g^2 C_A \int \frac{d^4 q}{(2\pi)^4} \left\{ \frac{\partial D_{\mu\nu}(p-q)}{\partial q_\sigma} \gamma_\mu S(q) \gamma_\nu - \frac{\partial}{\partial q_\sigma} [D_{\mu\nu}(p-q) \gamma_\mu S(q) \gamma_\nu] \right\} \\ = & -\frac{1}{2}g^2 C_A \int \frac{d^4 q}{(2\pi)^4} \frac{\partial D_{\mu\nu}(k)}{\partial k_\sigma} \gamma_\mu S(q) \gamma_\nu \\ = & -\frac{1}{2}g^2 C_A \int \frac{d^4 q}{(2\pi)^4} D_{\mu\alpha}(k) \Gamma_{\alpha\beta\sigma}^{3g}(k) D_{\beta\nu}(k) \gamma_\mu S(q) \gamma_\nu. \end{aligned} \quad (10)$$

It is apparent that the value of the above expression is not zero nonperturbatively due to the non-Abelian property[20]. Then we make clear the meaning of Eq. (8) and also prove that the claim made in Ref. [16] is incorrect.

Now an obvious question arises: whether there exists a truncation for the quark-gluon vertex to guarantee the the validity of the Slavnov-Taylor identity omitting the ghost effect? Recalling the above discussion, we can infer that such a truncation does exist. To make it realistic, we modify the ladder approximation of the quark-gluon vertex equation as

$$\begin{aligned} \Gamma_\sigma(p) = & i\gamma_\sigma - \left(C_F - \frac{C_A}{2} \right) g^2 \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(k) \gamma_\mu S(q) \Gamma_\sigma(q) S(q) \gamma_\nu \\ & + \frac{C_A}{2} g^2 \int \frac{d^4 q}{(2\pi)^4} D_{\mu\alpha}(k) \Gamma_{\alpha\beta\sigma}^{3g}(k) D_{\beta\nu}(k) \gamma_\mu S(q) \gamma_\nu. \end{aligned} \quad (11)$$

It is apparent that the second term in the right-hand side of the above equation is just the usual Abelian ladder approximation as shown in Eq. (2). An additional term is added in this expression which is related to the nonperturbative three-gluon vertex function. In this

sense, we should prove that the Slavnov-Taylor identity omitting the ghost effect is valid for Eq. (11). To reach such a point, we compare Eqs. (1) and (11). It is evident that Eq. (1) can be expressed in an equivalent form as

$$S(p)^{-1} = i\gamma \cdot p + m_0 + g^2(C_F - \frac{C_A}{2})\Sigma(p) + g^2\frac{C_A}{2}\Sigma(p), \quad (12)$$

where $\Sigma(p)$ is the quark self-energy in Eq. (7). Differentiating both sides of the above equation with respect to external momentum, one can get the general differential form of the quark equation

$$\begin{aligned} \frac{\partial S(p)^{-1}}{\partial p_\sigma} = & i\gamma_\sigma - \left(C_F - \frac{C_A}{2}\right) g^2 \int \frac{d^4 q}{(2\pi)^4} D_{\mu\nu}(k) \gamma_\mu S(q) \frac{\partial S(q)^{-1}}{\partial q_\sigma} S(q) \gamma_\nu \\ & + \frac{C_A}{2} g^2 \int \frac{d^4 q}{(2\pi)^4} D_{\mu\alpha}(k) \Gamma_{\alpha\beta\sigma}^{3g}(k) D_{\beta\nu}(k) \gamma_\mu S(q) \gamma_\nu, \end{aligned} \quad (13)$$

where the Slavnov-Taylor identity in Eq. (9) has been implemented for the third term in the right-hand side. Comparing Eq. (13) and Eq. (11), one can obtain the relation in Eq. (4). It shows apparently that the Slavnov-Taylor identity for the quark-gluon vertex omitting the ghost effects is valid for the modified ladder approximation with the modified vertex function in Eq. (11).

In summary, based on the Slavnov-Taylor identity for three-gluon vertex, we have proved that the claim made by Gogohia[16] is incorrect in this letter. Then the claim can not be used to say any thing about the internally inconsistent of the ladder approximation to QCD. Comparing with the argument provided in Ref. [17], our present statement is more physically and the reason for the contradiction between the Wark-Takahashi identity omitting ghost effects and the ladder approximation is represented more clearly and generally. Moreover, we provide a modification for the ladder approximation of the quark-gluon vertex which contains the usual Abelian term and an additional non-Abelian term. With such a modified ladder approximation to quark-gluon vertex and the rainbow approximation to quark equation we prove that the Slavnov-Taylor identity is valid for the quark-gluon vertex omitting the ghost effects.

This work was supported by the National Natural Science Foundation of China (NSFC) under contract Nos. 10425521 and 10575004, the Major State Basic Research Development Program under contract No. G2000077400, the research foundation of the Ministry of Education, China (MOEC), under contract No. 305001 and the Research Fund for the Doctoral Program of Higher Education of China under grant No 20040001010. One of the authors (YXL) thanks also the support of the Foundation for University Key Teacher by the MOEC.

References

- [1] W. Marciano and H. Pagels, Phys. Rep. **36** (1978) 137; H. Pagels, Phys. Rev. **D 15** (1971) 2991.
- [2] C. D. Roberts, and A. G. Williams, Prog. Part. Nucl. Phys. **33** (1994) 477.
- [3] C. D. Roberts, and S. M. Schmidt, Prog. Part. Nucl. Phys. **45** (2000) S1; and references therein.
- [4] R. Alkofer, and L. von Smekal, Phy. Rept. **353** (2001) 281.
- [5] P. Maris, and C. D. Roberts, Int. J. Mod. Phys. **E 12** (2003) 297; and references therein.
- [6] C. S. Fischer, R. Alkofer, W. Cassing, F. Llanes-Estrada, and P. Watson, Nucl. Phys. B (Proc. Suppl.) **153** (2006) 90.
- [7] R. Alkofer, P. Watson, and H. Weigel, Phys. Rev. **D 65** (2002) 094026.
- [8] A. Bender, D. Blaschke, Y. L. Kalinovsky, and C. D. Roberts, Phys. Rev. Lett. **77** (1996) 3724; T. Meissner, Phys. Lett. **B 405** (1997) 8; A. Bender, G. I. Poulis, C. D. Roberts, S. M. Schmidt, and A. W. Thomas, Phys. Lett. **B 431** (1998) 263; M. Harada, and A. Shibata, Phys. Rev. **D 59** (1999) 014010; T. Schaefer, and Schwenzer, Phys. Rev. **D 70** (2004) 054007; A. N. Mitra, and W. Y. P. Hwang, Eur. Phys. J. **C 39** (2005) 209.
- [9] C. S. Fischer, and R. Alkofer, Phys. Rev. **D 67** (2003) 094020; R. Alkofer, W. Detmold, C. S. Fischer, and P. Maris, Phys. Rev. **D 70** (2004) 014014.
- [10] R. T. Cahill, and C. D. Roberts, Phys. Rev. **D 32** (1985) 2419; M. A. Ivanov, Y. L. Kalinovsky, and C. D. Roberts, Phys. Rev. **D 60** (1999) 034018; P. Maris, and P. C. Tandy, Phys. Rev. **C 62** (2002) 055204; A. Bender, W. Detmold, A. W. Thomas, and C. D. Roberts, Phys. Rev. **C 65** (2002) 065203.
- [11] Y. X. Liu, D. F. Gao, and H. Guo, Nucl. Phys. **A 695** (2001) 353; Y. X. Liu, D. F. Gao, J. H. Zhou, and H. Guo, Nucl. Phys. **A 725**, 127 (2003); L. Chang, Y. X. Liu, and H. Guo, Nucl. Phys. **A 750** (2005) 324; L. Chang, Y. X. Liu, and H. Guo, Phys. Rev. **D 72** (2005) 094023.
- [12] H. T. Yang, H. S. Zong, J. L. Ping, and F. Wang, Phys. Lett. **B 557** (2003) 33; H. S. Zong, Y. M. Shi, W. M. Sun, and J. L. Ping, Phys. Rev. **C 73** (2006) 035206.
- [13] M. R. Pennington, QCD Down Under: Building Bridges, arXiv:hep-ph/0409156.

- [14] P. Watson, and R. Alkofer, Phys. Rev. Lett. **86** (2001) 5239; Z. G. Wang, and S. L. Wan, Phys. Lett. **B 536** (2002) 241; M. S. Bhagwat, M. A. Pichowsky, and P. C. Tandy, Phys. Rev. **D 67** (2003) 054019; P. Maris, A. Raya, C. D. Roberts, and S. M. Schmidt, Eur. Phys. J. **A 18** (2003) 231; R. Alkofer, M. Kloker, A. Krassnigg, and R. F. Wagenbrunn, Phys. Rev. Lett. **96** (2006) 022001.
- [15] W. Yuan, H. Chen, and Y.X. Liu, Phys. Lett. **B 637** (2006) 69 .
- [16] V. Gogohia, Phys. Lett. **B 611** (2005) 129.
- [17] H. S. Zong, W. M. Sun, Phys. Lett. **B 640** (2006) 196.
- [18] We use a Euclidean metric, with: $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$; $\gamma_\mu^\dagger = \gamma_\mu$; $a \cdot b = \sum_{i=1}^4 a_i b_i$; and $\text{tr}[\gamma_5 \gamma_\mu \gamma_\nu \gamma_\rho \gamma_\sigma] = -4 \epsilon_{\mu\nu\rho\sigma}$, $\epsilon_{1234} = 1$. For a spacelike vector k_μ , $k^2 > 0$. Landau gauge is used in our statements.
- [19] M. S. Bhagwat and P. C. Tandy, Phys. Rev. **D 70** (2004) 094039 .
- [20] A. I. Davydychev, P. Osland, and L. Saks, Phys. Rev. **D 63** (2001) 014022.